



# *Turfgrass and Environmental Research Online*

---

---

...Using Science to Benefit Golf



Ongoing research at Auburn University is being conducted to evaluate methods for converting bermudagrass fairways to zoysiagrass turf using various cultural and chemical methods. Results showed that dazomet and EPTC applied with siduron produced greater than 85% zoysiagrass cover by 15 weeks after establishment.

**Volume 8, Number 11**  
June 1, 2009

## PURPOSE

The purpose of *USGA Turfgrass and Environmental Research Online* is to effectively communicate the results of research projects funded under USGA's Turfgrass and Environmental Research Program to all who can benefit from such knowledge. Since 1983, the USGA has funded more than 350 projects at a cost of \$29 million. The private, non-profit research program provides funding opportunities to university faculty interested in working on environmental and turf management problems affecting golf courses. The outstanding playing conditions of today's golf courses are a direct result of ***using science to benefit golf.***

### Editor

Jeff Nus, Ph.D.  
1032 Rogers Place  
Lawrence, KS 66049  
jnus@usga.org  
(785) 832-2300  
(785) 832-9265 (fax)

### Research Director

Michael P. Kenna, Ph.D.  
P.O. Box 2227  
Stillwater, OK 74076  
mkenna@usga.org  
(405) 743-3900  
(405) 743-3910 (fax)

### USGA Turfgrass and Environmental Research Committee

Steve Smyers, *Co-chairman*  
Gene McClure, *Co-chairman*  
Julie Dionne, Ph.D.  
Ron Dodson  
Kimberly Erusha, Ph.D.  
Pete Grass, CGCS  
Ali Harivandi, Ph.D.  
Michael P. Kenna, Ph.D.  
Jeff Krans, Ph.D.  
James Moore  
Jeff Nus, Ph.D.  
Paul Rieke, Ph.D.  
James T. Snow  
Clark Throssell, Ph.D.  
Ned Tisserat, Ph.D.  
Scott Warnke, Ph.D.  
James Watson, Ph.D.  
Chris Williamson, Ph.D.

Permission to reproduce articles or material in the *USGA Turfgrass and Environmental Research Online* (ISSN 1541-0277) is granted to newspapers, periodicals, and educational institutions (unless specifically noted otherwise). Credit must be given to the author(s), the article title, and *USGA Turfgrass and Environmental Research Online* including issue and number. Copyright protection must be afforded. To reprint material in other media, written permission must be obtained from the USGA. In any case, neither articles nor other material may be copied or used for any advertising, promotion, or commercial purposes.

# Utilizing Soil Sterilants to Control Bermudagrass and Re-establish Zoysiagrass for Fairway Renovation

Mark C. Doroh, Joseph S. McElroy, and Robert H. Walker

## SUMMARY

Ongoing research at Auburn University is being conducted to evaluate methods for converting bermudagrass fairways to zoysiagrass turf using various cultural and chemical methods. Study results from summer 2008 indicate:

- Siduron (Tupersan) applied 24 lb/acre at zoysiagrass establishment resulted in poor bermudagrass control.
- Dazomet (Basamid) and EPTC (Eptam) applied with siduron or glyphosate (RoundUp Pro) resulted in less than 12% bermudagrass cover 15 weeks after establishment.
- Dazomet and EPTC applied with siduron produced greater than 85% zoysiagrass cover by 15 weeks after establishment.

Bermudagrass (*Cynodon spp.*) is extensively used throughout the southern United States for a variety of turfgrass purposes. Under high light intensity and warm temperatures, bermudagrass is a highly desirable turf species with many positive qualities. However, bermudagrass can also be difficult to control as a perennial weed when it contaminates other grass species.

A mixture of different grass species in the same area often leads to a poor turfgrass stand due to differences in color, texture, and growth-habit. A poor stand or mixture of grass species often necessitates control of the existing species prior to replanting of another grass. Bermudagrass especially presents a problem in zoysiagrass due to the limited availability of selective herbicides that do not excessively injure zoysiagrass (3, 5).

Renovating golf course fairways is an arduous task especially when converting from a mature stand of bermudagrass to another turf

species. Current renovation procedures of turf areas contaminated with bermudagrass are to fumigate or to treat with glyphosate and replant with the desired species (2, 10). Due to inconsistent control and turf regrowth, renovation with glyphosate has not proven to be a reliable method for eradicating unwanted bermudagrass (1, 4).

The alternative option for turfgrass renovation utilizes soil sterilants. Until 2005, methyl bromide was commonly used to kill established turf and fumigate the soil prior to replanting (6). However, with the phase-out process of methyl bromide already in motion, alternative soil fumigants have been assessed for their efficacy of disease and weed control (6, 7, 9, 10). Currently, little information is available about using alternative fumigants to eradicate bermudagrass and re-establish zoysiagrass for fairway renovation. Ongoing research at Auburn University is being conducted to evaluate methods for converting bermudagrass fairways to zoysiagrass using various herbicides and soil sterilants.



Bermudagrass can be difficult to control as a perennial weed when it contaminates other grass species.

MARK C. DOROH, Graduate Student; JOSEPH S. McELROY, Ph.D., Assistant Professor, Weed Science; and ROBERT H. WALKER, Ph.D., Turfgrass Weed Science; Agronomy and Soils Department, Auburn University, Auburn, AL.

Herbicide Treatments	Active Ingredient	Product Rate	Application Timing
RoundUp Pro	Glyphosate	1 gal/acre	5 Weeks Prior
Eptam	EPTC	1 gal/acre	3 Weeks Prior
RoundUp Pro	Glyphosate	1 gal/acre	5 Weeks Prior
Basamid	Dazomet	200 lb/acre	3 Weeks Prior
Eptam	EPTC	1 gal/acre	3 Weeks Prior
Basamid	Dazomet	200 lb/acre	3 Weeks Prior
Tupersan	Siduron	24 lb/ acre	At Establishment
RoundUp Pro	Glyphosate	1 gal/acre	5 Weeks Prior
Eptam	EPTC	1 gal/acre	3 Weeks Prior
Tupersan	Siduron	24 lb/acre	At Establishment
Basamid	Dazomet	200 lb/acre	3 Weeks Prior
Tupersan	Siduron	24 lb/acre	At Establishment
Non-treated	-----	-----	-----

**Table 1.** Herbicide treatments with related product rates and application timings.

### Materials and Methods

A fairway conversion study was initiated in May 2008 on 'Tifway 419' bermudagrass (*Cynodon dactylon* x *C. transvaalensis*) at the Auburn University Turfgrass Research Unit in Auburn, AL. Plot units measured 5 X 10 ft and were arranged in a randomized complete block design with four replications. Herbicide applications were applied at 3 mph with a CO<sub>2</sub> pressurized sprayer calibrated at 30 gal/acre with four Tee Jet XR8002US nozzles.

All areas were tilled and rolled prior to application of treatments. Soil sterilant treatments were incorporated with a second tillage. 'Zorro' zoysiagrass [*Zoysia matrella* (L.) Merr] was sprigged at a rate of 10 bushels/1000 ft<sup>2</sup>. A total of nine treatments were applied at either five weeks

prior, three weeks prior, or at establishment. Herbicide/sterilant treatments, rates, and application timings are listed in Table 1.

All plots received 0.5 lb N/1000 ft<sup>2</sup> every two weeks beginning six weeks after establishment. Tank mixtures of 2,4-D (0.95 lb/acre), dicamba (1.0 lb/acre), and halosulfuron-methyl (1.0 oz/acre) were used to control annual broadleaf and grassy weeds throughout the study. Treatments were visually rated at five, 10, and 15 weeks after establishment for percent zoysiagrass cover and bermudagrass contamination.

Bermudagrass and zoysiagrass cover were also subject to plant counts using a 48-inch (122 cm) dowel rod divided into 25 increments. The dowel rod was randomly placed twice within each plot. A species was counted upon intersection of an increment on the rod.




Figure 1. 'Zorro' zoysiagrass coverage 91 and 139 days after initial treatment.

## Results

Percent zoysiagrass and bermudagrass cover is presented in Figures 1 and 2. At 35 days after initial treatment, all treatments applied prior to establishment significantly reduced the amount of bermudagrass, yielding less than 15% ground cover. At 91 days after initial treatment (DAIT), the glyphosate + EPTC, glyphosate + dazomet, EPTC, EPTC + siduron, and dazomet + siduron treatments controlled bermudagrass well, yielding less than 15% ground cover. The dazomet alone and glyphosate alone treatments provided fair bermudagrass control at 31% and 34% ground cover, respectively.

Siduron applied at establishment controlled bermudagrass poorly, yielding 56% bermudagrass ground coverage. At 139 DAIT, the glyphosate + EPTC, glyphosate + dazomet, EPTC + siduron, and dazomet + siduron yielded less than 12% bermudagrass ground cover.

Treatments with good bermudagrass control yielded higher percent zoysiagrass cover than those with poor control. Dazomet and EPTC were the two dominant bermudagrass control treatments in this research. These two treatments applied with siduron or glyphosate provided equivalent control (less than 12% coverage).

EPTC equivalent control to dazomet is important economically. Dazomet can cost from \$800 to \$1400 per acre; whereas, EPTC estimated cost is approximately \$10 to \$20 per acre. However, comparing dazomet to EPTC applied alone, bermudagrass control was 25% and 75% respectively. Poor control using dazomet may be due to the low use rate (200 lb/acre) used in this research. Although previous studies with dazomet have used up to 350 lb/acre, we justify a lower use rate due to the high cost and the short time interval between application and replanting (three weeks). Trials will be replicated again in 2009 at the Auburn University Turfgrass Unit, Auburn,




Figure 2. 'Tifway' bermudagrass coverage 91 and 139 days after initial treatment.

AL. Herbicide and soil sterilant treatments will remain the same as those in 2008.

### Acknowledgements

The authors would like to thank the staff at the Auburn University Turfgrass Research Unit in Auburn, AL for their contribution to this work, as well as the USGA's Turfgrass and Environmental Research Program for providing funding for this research.

### Literature Cited

1. Boyd, J. 2000. Kill off bermudagrass with one less spraying: Replacing obsolete turf requires freedom from old bermudagrass. *Golf Course Management* 68:68-71. (TGIF Record 64769)
2. Cudney, D.W., C.L. Elmore, V.A. Gibeault, and J.S. Reints. 1997. Common bermudagrass (*Cynodon dactylon*) management in cool-season turfgrass. *Weed Technology* 11:478-483. (TGIF Record 40887)
3. Doernoeden, P. H. 1989. Bermudagrass suppression and zoysiagrass tolerance to fenoxaprop. Pages 285-290. In H. Takatoh (ed.). Proceedings of the Sixth International Turfgrass Research Conference. July 31-August 5, Tokyo, Japan. Japanese Soc. of Turfgrass Science, Tokyo. (TGIF Record 17106)
4. Johnson, B.J. 1988. Glyphosate and SC-0224 for bermudagrass (*Cynodon spp.*) cultivar control. *Weed Technology* 2:20-23. (TGIF Record 11928)
5. Johnson, B.J. 1992. Common bermudagrass (*Cynodon dactylon*) suppression in *Zoysia spp.* with herbicides. *Weed Technology* 6:813-819. (TGIF Record 29899)

6. Landschoot, P.J., and B.S. Park. 2004. Renovating putting greens without methyl bromide: A granular soil fumigant may be an option for putting green renovation if methyl bromide is banned in the near future. *Golf Course Management* 72(2):127-131. ([TGIF Record 127131](#))
7. Park, B.S., and P.J. Landschoot. 2003. A new product for fairway renovation: A granular soil fumigant may be the answer for superintendents looking at renovation projects. *Golf Course Management* 71(7):96-99. ([TGIF Record 87116](#))
8. McCarty, L.B. 2005. Best golf course management practices. 2nd ed. Pearson/Prentice Hall, Upper Saddle River, N.J. ([TGIF Record 101246](#))
9. Unruh, J.B., and B.J. Brecke. 2001. Seeking alternatives for methyl bromide: Nothing is ready as the ban nears, but some products show promise for fumigating prospective turf sites. *Golf Course Management* 69(3):65-71. ([TGIF Record 72411](#))
10. Unruh, J.B., B.J. Brecke, J.A. Dusky, and J.S. Godbehere. 2002. Fumigant alternatives for methyl bromide prior to turfgrass establishment. *Weed Technology* 16:379-387. ([TGIF Record 80617](#))