

Turfgrass and Environmental Research Online

Using Science to Benefit Golf



Researchers at the University of Massachusetts quantified dermal, airborne, and dislodgeable foliar residues (DFR) of three reduced-risk pesticides: azoxystrobin, carfentrazone-ethyl, and halofenozide. Dosimetry techniques were used to assess the exposure of volunteer golfers to these pesticides under worst-case scenarios during the play of 32 rounds of simulated golf. All exposures resulted in acquired doses significantly less than the reference doses (RfD) established for each pesticide by the USEPA Office of Pesticide Programs.

PURPOSE

The purpose of *USGA Turfgrass and Environmental Research Online* is to effectively communicate the results of research projects funded under USGA's Turfgrass and Environmental Research Program to all who can benefit from such knowledge. Since 1983, the USGA has funded more than 400 projects at a cost of \$31 million. The private, non-profit research program provides funding opportunities to university faculty interested in working on environmental and turf management problems affecting golf courses. The outstanding playing conditions of today's golf courses are a direct result of ***using science to benefit golf***.

Editor

Jeff Nus, Ph.D.
1032 Rogers Place
Lawrence, KS 66049
jnus@usga.org
(785) 832-2300
(785) 832-9265 (fax)

Research Director

Michael P. Kenna, Ph.D.
P.O. Box 2227
Stillwater, OK 74076
mkenna@usga.org
(405) 743-3900
(405) 743-3910 (fax)

USGA Turfgrass and Environmental Research Committee

Steve Smyers, *Chairman*
Gene McClure, *Co-chairman*
Ron Dodson
Kimberly Erusha, Ph.D.
Michael Fidanza, Ph.D.
Pete Grass, CGCS
Ali Harivandi, Ph.D.
Michael P. Kenna, Ph.D.
Jeff Krans, Ph.D.
James Moore
Jeff Nus, Ph.D.
Paul Rieke, Ph.D.
James T. Snow
Clark Throssell, Ph.D.
Scott Warnke, Ph.D.
Chris Williamson, Ph.D.

Permission to reproduce articles or material in the *USGA Turfgrass and Environmental Research Online* (ISSN 1541-0277) is granted to newspapers, periodicals, and educational institutions (unless specifically noted otherwise). Credit must be given to the author(s), the article title, and *USGA Turfgrass and Environmental Research Online* including issue and number. Copyright protection must be afforded. To reprint material in other media, written permission must be obtained from the USGA. In any case, neither articles nor other material may be copied or used for any advertising, promotion, or commercial purposes.

Utilizing Reduced-Risk Pesticides and IPM Strategies to Mitigate Golfer Exposure and Hazard

Jeffery J. Doherty and J. Marshall Clark

SUMMARY

Researchers at the University of Massachusetts quantified dermal, airborne, and dislodgeable foliar residues (DFR) of three reduced risk pesticides: azoxystrobin, carfentrazone-ethyl, and halofenozide. Dosimetry techniques were used to assess the exposure of volunteer golfers to these pesticides under worst-case scenarios during the play of 32 rounds of simulated golf. All exposures resulted in acquired doses significantly less than the reference doses (RfD) established for each pesticide by the USEPA Office of Pesticide Programs. The RfD is then used to calculate a hazard quotient (HQ) by dividing the acquired dose by the RfD. HQs ≤ 1.0 are indicative of a safe exposure. HQs >1.0 do not necessarily mean that the exposure is unsafe, only that the absence of adverse effects is less certain. Other findings include:

- Dermal absorption is the principle route of exposure to golfers following application of these reduced-risk compounds. Hands, lower legs, and upper socks were the main routes of exposure.
- Halofenozide exposure resulted in the highest combined hazard quotient (CHQ, sum of dermal and inhalation hazard quotients, 0.030) of the reduced risk pesticides tested following application at its highest labeled rate with 0.25" of post-application irrigation. It should be noted that halofenozide has recently lost its reduced-risk pesticide designation. The CHQ calculation for halofenozide is only about half of that determined by dosimetry for chlorpyrifos (0.059). In this case, the application of halofenozide is considered safe, but does not greatly reduce golfer hazard.
- The CHQ established for the fungicide azoxystrobin (0.0039) was about an order of magnitude less than that determined for the fungicide chlorothalonil (0.043). The CHQ determined for the herbicide carfentrazone-ethyl (0.00014) was ~ 700-fold less than that determined for the herbicide MCPP (0.1).
- In the cases examined, the use of reduced-risk pesticides can lessen golfer exposure and hazard. The application of azoxystrobin brought about an ~ 10-fold reduction in CHQ compared to chlorothalonil. The use of carfentrazone-ethyl produced an ~ 700 fold reduction in CHQ versus the use of MCPP, and the application of halofenozide led to a ~ 2-fold reduction in HQ compared to the use of chlorpyrifos.
- A 1-hour reentry interval generally reduces exposure.

Pesticide exposure following application to turfgrass is of concern to turf managers, golfers, and regulators. Although most consumers are not well informed about pesticide use on golf courses, well-publicized examples of environmental contamination from agrochemicals have caused many individuals to develop negative attitudes toward golf course management practices. These concerns arise from the variety, amount, and frequency of pesticide use on turf, the amount of time adults and especially children and infants spend on turfgrass, and the potential for unsafe levels of exposure that may occur in these situations. There has been a great deal of focus on the exposures of

mixers, loaders, and applicators to pesticides. The potential for exposure, however, exists for all who re-enter pesticide-treated turf areas.

Previous Studies

Our previous results indicate that the overall "golfer" exposure received following full-course and full-rate applications of chlorpyrifos, carbaryl, and cyfluthrin, under worst case scenarios, are substantially below USEPA RfD values, indicating safe exposures (CHQ ≤ 1.0 ; combined hazard quotient; 4). The initial phase of this project focused on studying the well-characterized conventional insecticides chlorpyrifos and carbaryl, as well as a more modern insecticide, cyfluthrin. Insecticides were chosen in the belief that these neurotoxins would present the most

JEFFERY J. DOHERTY, Graduate Student; and J. MARSHALL CLARK, Ph.D., Director, Massachusetts Pesticide Analysis Laboratory Veterinary and Animal Science Department, University of Massachusetts, Amherst MA

USGA Turfgrass and Environmental Research Online 10(15):1-18.

TGIF Record Number: 187985




Figure 1. The transfer factor is the difference between the dislodgeable foliar residues (DFRs) on the turfgrass and the amounts determined on the dosimetry suits worn by a volunteer golfer simulating a round of golf. By segmenting the suits the routes of exposure can be determined.

hazard to golfers. All insecticides received 0.25 inches of post-application irrigation. Putnam and Clark (11) determined CHQs for golfer exposure to chlorpyrifos (0.059), carbaryl (0.014), and cyfluthrin (0.004) (4). All CHQs were well below 1.0, indicating safe exposures. It was also determined that the lower body was the major route of exposure.

The next phase of the project looked at conventional herbicides (2,4-D, MCPP), a commonly used fungicide (chlorothalonil), and a more modern insecticide (imidacloprid). As with the other insecticides, imidacloprid received 0.25 inches of post-application irrigation. Herbicides and fungicides did not receive post-application irrigation. This study determined that the CHQs for 2,4-D (0.011), MCPP (0.55), imidacloprid (0.0097), and chlorothalonil (0.043) were also well below USEPA HQ values. Golfer exposure to MCPP, a herbicide, resulted in the highest CHQ generated in this project. Without post-application irrigation (2,4-D, MCPP, chlorothalonil), the primary route

of exposure was through the hands. With post-application irrigation, imidacloprid exposure was determined to be through the lower body, much like the exposure scenarios of the previously studied insecticides.

Justification of Worst Case Scenarios

In all these studies, we consider these exposure estimates to be worst-case scenarios for the following reasons:

- All pesticides were applied at maximum-labeled rates.
- Pesticide applications were made to the entire “course”, including tees, greens, collars, and fairways.
- Volunteer golfers spent the entire four-hour round of golf on the treated turf.
- USEPA chronic RfDs, which assume no harmful effects following daily exposure over a 70-year lifespan, were used to generate the HQs.

In this study, we measured environmental residues (DFR, dislodgeable foliar residues) available for transfer from treated turf, and residues on dosimeters (whole-body cotton suits and personal air samplers) to measure the amount of pesticide on the body surface available for dermal penetration and that in the personal breathing space available for inhalation, respectively. This information provides a novel and complete database for the assessment of pesticide exposure from treated turf.

Our previous work focused on strategies for mitigating golfer exposure to turfgrass pesticides such as half-course applications, application only to tees and greens, and use of post-application irrigation when appropriate. These strategies were found to reduce golfer exposure and hazard. Reduced-risk pesticides, which typically have reduced mammalian toxicity, may be yet another strategy to reduce golfer exposure and hazard. For this reason, we chose to examine the effect of the use of reduced-risk pesticides on golfer exposure and hazard.

For a pesticide to be considered reduced-risk, it must meet one or more of the following criteria:

Human Health Effects

- very low mammalian toxicity
- toxicity generally lower than alternatives (10-100x)
- displaces chemicals that pose potential human health concerns (e.g. organophosphates, or probable carcinogens)
- reduces exposure to mixers, loaders, applicators, and re-entry workers

Non-Target Organism Effects (birds and insect)

- very low toxicity to birds and beneficial insects
- very low toxicity to honeybees
- significantly less toxic to birds than alternatives
- not harmful to beneficial insects; highly selective pest impacts

Non-Target Organism effects (fish)

- very low toxicity to fish
- less toxicity/risk to fish than alternatives
- potential toxicity/risk to fish mitigatable
- similar toxicity to fish as alternatives, but significantly less exposure



Figure 2. A Toro Multipro 1200 boom sprayer fitted with eleven VisiFlo® flat spray tips (TeeJet® Technologies) was used for all applications.

Pesticide (rate)	Post-application Irrigation	Exposure (h)	Field Trials
Azoxystrobin Heritage (0.2 oz. a.i./1,000 ft ²)	None	4	3
Carfentrazone-ethyl Quicksilver T&O (0.036 oz. a.i./1,000 ft ²)	None	4	2
Halofenozide Mach II 2SC (0.65 oz. a.i./1,000 ft ²)	0.25"	4	3

Table 1. Summary of reduced-risk pesticides tested, application rates and conditions, and duration and frequency of exposures.

Groundwater Issues

- low potential for groundwater contamination
- low drift, runoff potential
- runoff mitigatable
- lower use rates than alternatives
- low pest resistance potential (e.g. new mode of action)
- highly compatible with IPM
- effective at very low use rates

Routes of Pesticide Exposure

Pesticide exposure to golfers is principally through the dermal contact with dislodgeable foliar residues on treated turfgrass foliage. The dense canopy of the turfgrass system is likely to intercept a greater percentage of applied pesticide than typical agricultural systems where much of the applied pesticide reaches the soil surface. Therefore, dermal exposure may be significant. Turfgrass plants also have a waxy layer on their surface and also produce an organically rich thatch layer. These factors are expected to compete with the transfer of pesticides to golfers.

Inhalation is a second possible route of golfer exposure. These reduced-risk pesticides are relatively non-volatile, making airborne exposure to these compounds less of a concern. This exposure

route is still toxicologically relevant, however, because inhalation exposure results in a high level of absorption. The lungs highly vasculated circulatory system allows for extremely rapid transfer of volatiles and small particulates contained in inhaled air to the bloodstream and widespread distribution throughout the body.

Oral exposure is the third possible route, but is generally considered the least important in a golfer exposure scenario. This type of exposure would primarily be through hand-to-mouth contact. Evidence has shown that golf balls, tees, etc. do not transfer significant amounts of pesticides to golfers (3).

Material and Methods

Field Study Site

All experiments were performed at the University of Massachusetts Joseph Troll Turfgrass Research Center in South Deerfield MA. A 130 X 30-yard plot consisting of 'Penncross' creeping bentgrass was used for the concurrent collection of DFR and dosimetry samples. This plot was maintained as a golf course fairway, mowed three times a week at 0.5-inch, and irrigated as needed to prevent drought stress.



Figure 3. Each experiment utilized four volunteers wearing dosimetry suits simulating the play of an 18-hole round of golf over a period of four hours. The dosimetry group of volunteers wore white, 100% cotton, long johns, white cotton gloves, and veils attached to the back of their hats

Additionally, the plot was fertilized with 3 lbs N/year in three 1-lb applications, and treated with fungicides not used in the current study to control fungal diseases [dollar spot (*S. homeocarpa*), brown patch (*R. solani*), and pythium blight (*Pythium* spp.)] as needed.

Pesticide Applications

A Toro Multipro 1200 boom sprayer fitted with eleven VisiFlo® flat spray tips (TeeJet® Technologies) was used for all applications (Figure 2). All treatments with MachII 2SC® (halofenozide, Syngenta) were followed by 0.64 cm (0.25”) of irrigation immediately after application.

Heritage® (50% azoxystrobin, Dow

Agrosciences) was applied at a rate of 0.2 oz. a.i./1,000 ft². Heritage® was not applied at the maximum label rate of 0.35 oz. a.i./1,000 ft² used for snow mold control because these applications take place shortly before snow cover when the golf season is already over. This would, therefore, present an unrealistic exposure scenario. Briefly, 16 oz. of formulated product was mixed with 80 gallons of water and applied at a rate of 2 gallons/1,000 ft².

Quicksilver T & O Herbicide® (21.3.0% carfentrazone-ethyl, FMC Corp.) was applied at the USEPA maximum label rate of 0.036 oz. a.i./1,000 ft². For the Quicksilver application, 6.16 oz. of formulated product was mixed with 80 gallons of water and applied at a rate of 2 gallons/1,000 ft².




Figure 4. Inhalation exposure to the dosimetry group was measured utilizing personal air samplers (Aircheck52, SKC Inc., Eighty Four PA) equipped with OVS air sampling tubes (SKC Inc.) attached to the front collar of each volunteer near the breathing zone.

Mach II 2SC® (22.3% halofenozide, Syngenta) was applied at the USEPA maximum label rate of 0.65 oz. a.i./1,000 ft². For the Mach II application, 7.25 qt. (116 fluid oz.) of formulated product was mixed into 80 gallons of water and applied at a rate of 2 gallons/1,000 ft².

Golfer Activities and Exposure Scenarios

Azoxystrobin, carfentrazone-ethyl, and halofenozide were applied to a rectangular bent-grass plot maintained as a fairway. Exposure to volunteers simulating the play of a round of golf was determined by a dosimetry study. Concurrent with this study, DFR samples were collected from the same plot. Each experiment utilized four volunteers wearing dosimetry suits simulating the play of an 18-hole round of golf over a period of four hours.

The round of golf was standardized in an attempt to ensure consistent behaviors among different volunteers. The template was based on all volunteers playing bogey golf (par plus one stroke per hole). Each player hit the ball 90 times, addi-

tionally taking 90 practice swings, while walking the 6,500-yard layout of a local course. Clubs were rotated in an appropriate manner, balls were teed up on tee shots, divots were replaced, and clubs were wiped clean between shots using a golf towel. Each simulated round of golf commenced either one-hour after application (azoxystrobin, carfentrazone-ethyl) or one hour after the end of post-application irrigation (halofenozide).

Dislodgeable Foliar Residues (DFRs)

These experiments were carried out on the same plot concurrent with the exposure study. DFRs were determined using the Outdoor Residential Task Force recommended California roller (CA roller, Figure 5) (3) Three 7' x 12' sections of the study site were cordoned off to prevent foot traffic by the study participants. One 2' x 3' DFR sample was collected from each section at 0.25, 1, 2, and 5 hours after pesticide application. Sections that were sampled had their perimeters marked using turf paint to prevent re-sampling of an area.

Dosimetry

Exposure of volunteers simulating the play of an 18-hole round of golf was determined by dosimetry. Exposure to individual volunteers has been evaluated following 32 rounds of golf derived from 8 pesticide applications to the study site (Table 1). The dosimetry group of volunteers wore white, 100% cotton, long johns (Indera Mills Co., Yadkinville, NC), white cotton gloves, and veils attached to the back of their hats (Figure 3), which served as passive collection media for pesticide residues from treated turfgrass (5, 6).

Inhalation exposure to the dosimetry group was measured utilizing personal air samplers (Aircheck52, SKC Inc., Eighty Four PA) equipped with OVS air sampling tubes (SKC Inc.) attached to the front collar of each volunteer near the breathing zone (Figure 4)(5, 7). These sampling tubes trap three types of pesticide residues: those sorbed onto particulates (e.g., dust) and those found in aerosols are trapped by a quartz fil-

ter, and pesticide vapors, which are trapped on XAD-2 resin in the tubes. Air samples were collected at 2.0 L/min. To estimate total airborne dose of the pesticide, the residues found in the sampling tubes were multiplied by 10.5 to arrive at the dose correlated with a moderate breathing rate of 21 L/min.

Volunteers were from the Department of Veterinary and Animal Science at the University of Massachusetts, Amherst. A protocol that describes the study and the protected rights of the volunteers was approved by the Human Subjects Review Committee at UMASS (OGCA# 107-0889). The approved protocol, including an informed consent form, was reviewed with potential participants at an orientation meeting prior to their participation.

All pesticide analysis was carried out at the Massachusetts Pesticide Analysis Laboratory (MPAL), a USEPA/MA Department of Agricultural Resources (DAR)-supported pesticide analytical laboratory using standard protocols and QA/QC procedures.

Results

Airborne and DFRs were determined for a period of 4.75 hours following pesticide application, or 4.75 hours after the end of post-application irrigation in the case of halofenozide. Foliar dislodgeable residues were quantified using the CA roller device, and airborne residues were determined by personal air samplers. These environmental samples were collected concurrent with the dosimetry exposure study on the same plot.

Azoxystrobin DFRs

In agreement with previous research on chlorpyrifos and carbaryl (4), azoxystrobin residues declined rapidly in the first hour following application, then exhibited a slower decline over the next four hours (Figure 6). Azoxystrobin DFRs declined from 0.60 $\mu\text{g}/\text{cm}^2$ (± 0.04 S.E) at 0.25 hours to 0.34 $\mu\text{g}/\text{cm}^2$ (± 0.03) at one hour (43% reduction) and then to 0.02 $\mu\text{g}/\text{cm}^2$ (± 0.005) in the next four hours for a total reduction



Figure 5. Foliar dislodgeable residues were quantified using the California roller device shown above.

of 97% in available residues over the 4.75 hours of the experiment. This dissipation pattern indicates that it is appropriate for DFRs to be averaged over a four-hour period, our standardized playing time for an 18-hole round of golf, for use in exposure estimates (1). Using this approach, a mean DFR of 0.146 (± 0.015) μg azoxystrobin/ cm^2 was calculated for the time course of the experiment (hours 1-5) following application at a rate of 0.2 oz. a.i./1,000 ft^2 .

Carfentrazone-ethyl DFRs

Dissipation of DFRs of carfentrazone-ethyl followed a similar pattern (Figure 7). DFRs of carfentrazone-ethyl declined from 0.00082 $\mu\text{g}/\text{cm}^2$ (± 0.000011) to 0.00052 $\mu\text{g}/\text{cm}^2$ (± 0.000016) at one hour (a 37% reduction), and then to 0.00026 $\mu\text{g}/\text{cm}^2$ (± 0.000030) in the next four hours, resulting in an overall reduction of 68% in available residues. A mean DFR was calculated as 0.0005 (± 0.000014) μg carfentrazone-ethyl/ cm^2 for the time course of the experiment (hours 1-5) following application at a rate of 0.035 oz. a.i./1,000 ft^2 .

Halofenozide DFRs

The dissipation pattern of halofenozide was not consistent with previous results (Figure 8).




Figure 6. Availability of azoxystrobin dislodgeable foliar residues (DFR) over the first five hours following application.

Halofenozide residues did not significantly decline in the first hour following application. Dislodgeable foliar halofenozide residues only declined from 0.295 µg halofenozide/cm² (± 0.072) at 0.25 hours to 0.276 µg/cm² (± 0.078) at one hour and this reduction was not significantly different. Residues then declined to 0.023 µg/cm² (± 0.01) over the next four hours for an overall reduction of 92% of available residues. A mean DFR was determined to be 0.115 µg/cm² (± 0.022) for the time course of the experiment (hours 1-5) following application at a rate of 0.65 oz. a.i./1,000 ft². It should be noted that halofenozide is a systemic insecticide taken up by the roots and has a relatively high water solubility of 12.3 ppm (7).

The dissipation pattern for 2,4-D, MCPP, chlorothalonil, and imidacloprid were consistent with the results for azoxystrobin, carfentrazone-

ethyl, and the insecticides studied earlier (chlorpyrifos, carbaryl, and cyfluthrin). The dissipation pattern of halofenozide is unique to all the pesticides we have studied to date.

Determination of Exposure by Dosimetry

The total residues of azoxystrobin, carfentrazone-ethyl, and halofenozide collected on whole-body dosimeters and personal air samplers during the simulated play of an 18-hole round of golf are given in Table 2. An average of ~ 800 µg of azoxystrobin was collected on the whole body dosimeters following three applications of Heritage® at a rate of 0.2 oz. a.i./1,000 ft² (Table 2). After adjusting for a moderate breathing rate (21 L/min.), personal air samplers worn by the dosimetry group collected an average of 0.156 µg of azoxystrobin. As shown previously (4), the principle route of exposure to golfers was dermal




Figure 7. Availability of carfentrazone-ethyl dislodgeable foliar residues (DFR) over the first five hours following application.

following chlorpyrifos application. In the case of azoxystrobin, similar results were found. Dermal residues account for >99.9% of all transferred residues.

DFR Distribution

The distribution of azoxystrobin on the various dosimeter segments is shown in Figure 9. The hands were the primary route of exposure to axoxystrobin, accounting for ~ 55% of the total residues on the dosimeters, while the upper sock accounted for an additional ~ 27%. Together, the hands and lower sock account for ~ 82% of the total pesticide residues transferred to the dosimeter. The remaining dosimeter segments (veil, upper and lower arm, torso, pant, and lower leg) accounted for ~ 18% of the total residues combined.

Total residues of carfentrazone-ethyl found on the dosimeters following two applica-

tions of Quicksilver T&O® at a rate of 0.036 oz. a.i./1,000 ft² were determined to be ~ 2.9 µg (Table 2). No carfentrazone-ethyl was detected in the personal air samplers, so these samples were assumed to contain residues equal to half the limit of detection for these samples. After adjusting for a moderate breathing rate, the estimated airborne exposure to carfentrazone-ethyl was 0.006 µg. Dermal residues accounted for 99% of all transferred residues.

Carfentrazone-ethyl distribution on the dosimeters is shown in Figure 10. Again, the hands (~ 28% of transferable residues) were the primary route of exposure, but the pants held the next highest percentage of residues at ~ 22%. These two segments account for ~ 50% of all dermal residues of carfentrazone-ethyl. Overall, carfentrazone-ethyl was more widely distributed on the dosimeter than the other two compounds in this study. It is important to note that roughly half the dosimeter segments analyzed for carfentra-




Figure 8. Availability of halofenozide dislodgeable foliar residues (DFR) over the first five hours following application.

zone-ethyl were negative, including all the veils. Because of this, many suit segments were assigned a residue concentration equal to half the limit of detection ($0.1\mu\text{g}/\text{segment}$). All dosimeter suit segments from the azoxystrobin and halofenozide analyses were positive for those pesticides (above the limit of detection).

An average of approximately $940\mu\text{g}$ of halofenozide was found on the whole body dosimeters following three applications of MachII SC® at a rate of $0.65\text{ oz. a.i./1,000 ft}^2$ (Table 2). Halofenozide was the only one of the three pesticides that received post-application irrigation. Personal air samplers worn by the dosimetry group collected an average of $0.095\mu\text{g}$ of halofenozide, assuming a moderate breathing rate, over the four-hour simulated round of golf. Again, the principle route of exposure was dermal. With halofenozide, $> 99.99\%$ of the transferred residues were dermal.

As with azoxystrobin, the hands were the prin-

ciple route of exposure to halofenozide, accounting for $\sim 54\%$ of the total dermal exposure (Figure 11). The lower leg ($\sim 17\%$) and the upper sock ($\sim 13\%$) account for an additional $\sim 30\%$ of the dosimeter residues. The rest of the dosimeter (veil, upper arm, lower arm, torso, and pant) contained $\sim 16\%$ of the total dermal residues combined.

Calculation of Dermal, Airborne, and Combined Hazard Quotients

Dosimetry data alone can be used to directly calculate a dermal hazard quotient (DHQ). Using azoxystrobin as an example, we have determined the total residues of azoxystrobin on the dosimeter to be $805\mu\text{g}$ following application of Heritage® at $0.2\text{ oz. a.i./1,000 ft}^2$. Using USEPA or EU Pesticide Properties Database established values for dermal penetration (6.1% for azoxystrobin), USEPA values for the weight of

an average person (70kg), and the chronic RfD (0.18 mg/kg/d for azoxystrobin), the DHQ for azoxystrobin is determined using Equation 1.

Equation 1

$$\text{DHQ} = \frac{805 \mu\text{g azoxystrobin} \times 6.1\% \text{ penetration factor}/70 \text{ kg}}{180 \mu\text{g/kg/day (USEPA Azoxystrobin RfD)}}$$

DHQ = 0.0039

Using the same approach, we calculated a DHQ for carfentrazone-ethyl of 0.00014 (Table 2). Halofenozide has no established EPA dermal penetration factor, so a default value of 20% was used. Since halofenozide is not registered for any food crop uses, there is no USEPA established RfD either. Using standard EPA methodology, a chronic RfD of 38 µg/kg/day has been calculated (8). Using this estimated penetration factor and RfD for halofenozide, a dermal hazard quotient of 0.030 is calculated.

The residues collected on the personal air samplers can also be used to calculate an inhalation hazard quotients (IHQ). Again using azoxystrobin as an example, the average residues found on the air samplers following application of Heritage® at 0.2 oz. a.i./1,000 ft² was 0.156 µg. The IHQ can be calculated in a manner similar to the DHQ.

Equation 2

$$\text{IHQ} = \frac{0.156 \mu\text{g}/70 \text{ kg}}{180 \mu\text{g/kg/day (USEPA Azoxystrobin RfD)}}$$

IHQ = 0.000012

Using the same approach, we calculated IHQs for carfentrazone-ethyl and halofenozide (Table 2). No carfentrazone-ethyl residues were ever detected in the personal air samplers. We therefore assigned these residues a value half that of the limit if detection to arrive at an IHQ of 0.0000026 for carfentrazone-ethyl. The calculation for the IHQ of halofenozide resulted in an IHQ of 0.000037.

These hazard quotients can then simply be summed to determine a combined hazard quotient (CHQ) as follows:

Azoxystrobin Combined Hazard Quotient (CHC)

0.0039 DHQ + 0.000012 IHQ = 0.003912

Carfentrazone-ethyl Combined Hazard Quotient (CHC)

0.00014 DHQ + 0.0000026 IHQ = 0.0001426

Halofenozide Combined Hazard Quotient (CHC)

0.030 DHQ + 0.000037 IHQ = 0.030037

Application Rate	Mean Suit Residues (µg)	Mean Airborne Residues (µg)	DHQ	IHQ	Combined HQ
Azoxystrobin (0.2 oz a.i./1,000 ft ²)	805	0.156	0.0039	0.000012	0.003912
Carfentrazone (0.036 oz. a.i./1,000 ft ²)	2.86	0.0055	0.00014	0.0000026	0.0014
Halofenozide (0.65 oz. a.i./1,000 ft ²)	944	0.095	0.030	0.0014	0.03014

Table 2. Dermal hazard quotients (DHQ), inhalation hazard quotients (IHQ) and overall hazard quotients (Overall HQ) calculated from dosimetry data for the three pesticides studied. No detectable carfentrazone-ethyl residues were found. The value given is determined as half the limit of detection for the sample.




Figure 9. The distribution of azoxystrobin on whole body dosimeters.

These data clearly show that airborne residues of the three reduced-risk pesticides examined contribute very little to the CHQ of each of the three pesticides. It should also be noted that all three CHQs are significantly less than 1.0, indicating a wide safety margin. The CHQs of azoxystrobin, carfentrazone-ethyl, and halofenozide were 250-, 7000-, and 14-fold below the current USEPA level of concern respectively.

The use of the chronic RfD also adds to this safety margin. A chronic RfD is one that can be received daily over a 70-year lifespan with no adverse effects. Using the same experimental design, we have previously calculated CHQs for 2,4-D (0.011), MCPP (0.55), chlorothalonil (0.043), and imidacloprid (0.01) using dosimetry.

Transfer Factors

Using the dermal dose derived from dosimetry

and the DFRs determined from the environmental residues, we can calculate a transfer factor (TF).

Equation 3

$$TF(\text{cm}^2/\text{h}) = \text{dermal exposure } (\mu\text{g}) / \text{DFR } (\mu\text{g}/\text{cm}^2)/4\text{h}$$

Using azoxystrobin as an example, we can calculate a TF as follows:

$$\text{Azoxystrobin TF} = 805 \mu\text{g} / 0.146 \mu\text{g}/\text{cm}^2/4\text{h}$$

$$\text{Azoxystrobin TF} = \sim 1,380$$

Employing the same methodology, transfer factors have been calculated for carfentrazone-ethyl (1,440) and halofenozide (2,050). Transfer factors have also been calculated for 2,4-D (570), MCPP (1,300), chlorothalonil (1,150), and imidacloprid (3,300).

Knowing the DFR and the TF allows an

absorbed dose to be established solely from these two values as well as the dermal penetration of the pesticide. The estimated absorbed dose is determined as follows where AD = absorbed dose and DP = dermal penetration.

Equation 4

$$AD = DFR (\mu\text{g}/\text{cm}^2) \times TF (\text{cm}^2/\text{h}) \times DP \times 4\text{h}/70\text{kg}$$

Utilizing this equation allows an absorbed dose to be estimated without the need for time-consuming and expensive golfer exposure studies using human volunteers.

Discussion

Dermal exposure estimates based on full-course applications at the maximum label rates with a one-hour re-entry interval for azoxystrobin, carfentrazone-ethyl, and halofenozide resulted in USEPA HQ values substantially below 1.0, indicating safe exposures. Previously, Putnam and Clark (10) showed good agreement between HQs generated by dosimetry and biomonitoring data. This finding indicates that in the absence of voided urinary metabolites, dosimetry can be an adequate surrogate for biomonitoring if transfer and penetration factors are known. The measurement of dosimetry residues combined with airborne residues and DFRs provide a complete and accurate account of pesticide availability for golfer exposure. Pesticide regulators have a critical need for this type of data.

The current study evaluated whether reduced-risk pesticides could lessen golfer hazard. Reduced-risk pesticides typically have lower mammalian toxicities and lower application rates in addition to their other environmental benefits compared to conventional pesticides. These factors indicate that reduced-risk pesticides should lessen golfer hazard. This proved to be the case for azoxystrobin and carfentrazone-ethyl compared to chlorothalonil and MCP, respectively. Halofenozide on the other hand, did not result in lower HQs compared to chlorpyrifos, and was

actually significantly higher than that found for cyfluthrin. Nevertheless, all pesticide exposures resulted in HQs less than 1.0.

HQ determination depends primarily on the RfD. The USEPA determines the chronic (daily exposure over a lifetime) RfD by first defining the No Observed Adverse Effect Level (NOAEL), which is the highest dose in a chronic toxicological study that induces no adverse effect in the animal. This NOAEL is then divided by an uncertainty factor (UF) taking into account differences between species (10X) and hazard to vulnerable populations such as children (10X). The typical UF is 100X. Only in cases where there is sufficient human data, such as with chlorpyrifos, is the UF = 10. Only three pesticides have UFs of 10 (aldicarb, chlorpyrifos, and malathion), and only acephate (UF = 30) and pirimiphos-methyl (UF = 25) are also below 100.

The UF of 10 is due to the fact that differences between species no longer has to be accounted for in the RfD calculation. Such low uncertainty factors are extremely rare. If the USEPA determines that there is insufficient toxicological data such as use of too few species, they will introduce another 10X safety factor, making a UF of 1,000. In cases where there is extremely limited toxicological data, they may assign further safety factors such as with MCP, which has an uncertainty factor of 3,000 due to an additional 3X safety factor.

The azoxystrobin RfD is calculated based on a NOAEL of 18.2 mg/kg/day from a chronic rat feeding study as follows using equation 5:

Equation 5

$$RfD = \frac{NOAEL}{UF}$$

$$\text{Azoxystrobin RfD} = \frac{18.2 \text{ mg/kg/day}}{100} = 182 \mu\text{g/kg/day}$$

While the RfD is the driving force behind HQ determination, factors that influence exposure such as application rate, water solubility, and Koc can also influence the HQ (Table 3).

The HQs of two of the reduced-risk pesti-




Figure 10. The distribution of carfentrazone-ethyl on whole body dosimeters. Hands and pants received almost ½ the dose. * Veils (head) had no detectable residues. Pants were the only segment with all samples being positive for carfentrazone-ethyl. 25% to 75% of the individual samples for all other segments had no detectable residues.

cides were significantly less than comparable, commonly used turfgrass pesticides. Azoxystrobin exposure following a simulated round of golf commencing 1 hour after application resulted in a HQ >250-fold below current levels of concern. This compares favorably to the HQ for chlorothalonil of 23-fold below the level of concern following the same exposure scenario. The CHQ generated for carfentrazone-ethyl was >7,000-fold below the USEPA’s level of concern. This hazard is significantly lower than that found for MCP, which was 10-fold below this level.

The CHQ resulting from exposure of volunteer golfers to halofenozide 1 hour after post-application irrigation presents a different story. This CHQ was 33-fold below the USEPA’s current level of concern. This hazard is about twice that

generated for chlorpyrifos in previous work of 17-fold below the current level of concern (11), and is much higher than that determined for cyfluthrin (250-fold below).

The generally higher costs of reduced-risk pesticides may give turfgrass managers pause when considering their use. We have demonstrated another benefit in their use by reducing golfer hazard, in addition to their utility in an IPM program. The high margins of safety for azoxystrobin and carfentrazone-ethyl should ensure that these chemicals remain registered for turf, and may help EPA “fast track” approval of similar pesticides for use on turf.

The CHQ generated for the fungicide azoxystrobin (0.0039) was about an order of magnitude less than that determined for chlorothalonil

Pesticide	Application Rate (oz. a.i./1,000 ft ²)	Water Solubility (mg/L)	Koc	Transfer Factor	Uncertainty Factor	Dermal Penetration (%)	RfD (µg/kg/d)	CHQ
Azoxystrobin	0.2	6	482	1,378	100	6.1%	180	0.0039
Chlorothalonil	4.16	0.81	5,000	1,150	100	0.15%	15	0.040
Carfentrazone	0.036	12	866	1,436	100	10%	30	0.00014
MCPPP	0.13	860	20-43	1,300	3,000	16.3%	1.0	0.55
Halofenozide	0.65	250	250	2,051	1,000	20%	38	0.070
Chlorpyrifos	0.38	1.4	9,930	6,400	10	9.6%	5	0.059
Cyfluthrin	0.053	0.007	64,300	1,075	100	5%	25	0.004

Table 3. A summary of the factors influencing the determination of hazard quotients. Koc = water/organic carbon partition coefficient, RfD = USEPA chronic reference dose, CHQ = combined hazard quotient.

(0.043), a conventional fungicide. This finding roughly parallels 12-fold difference in RfD between azoxystrobin (182 µg/kg/day) and chlorothalonil (15 µg/kg/day).

The RfD does not explain all the difference, however, as the application rate for chlorothalonil was approximately 20 times that of azoxystrobin, and the transfer factors were similar, meaning that roughly the same ratio of available residues were transferred to the dosimeter. The rest of the difference may be explained by the extremely low dermal penetration rate for chlorothalonil (0.15%) compared to azoxystrobin (6.1%).

Exposure to the herbicide carfentrazone-ethyl following simulated golf resulted in a CHQ of 0.00014. The carfentrazone-ethyl CHQ was almost 4,000-fold below that determined for MCPPP (CHQ=0.55). Some of this higher CHQ can be explained by the RfD for MCPPP, which is 30-fold lower than that for carfentrazone-ethyl. All of this difference in RfD can be explained by the UF of 3,000 assigned to MCPPP. This is 30-fold higher than the UF associated with carfentrazone-ethyl. Additionally, the application rate for carfentrazone-ethyl was ~ 3.6-fold lower than for MCPPP.

It is clear from comparing the CHQs of these fungicides and herbicides that the use of reduced-risk compounds can reduce golfer hazard. This finding is important for turfgrass managers as it adds another positive factor for the use of reduced-risk compounds in addition to their utili-

ty in IPM programs due to their novel modes of action and their enhanced environmental safety.

The above results were in contrast to those found for halofenozide, which recently lost its USEPA reduced-risk designation. The CHQ determined for halofenozide (CHQ = 0.070) was close to that determined for chlorpyrifos (CHQ = 0.059) (4), and was ~ 18-fold lower than that determined for cyfluthrin (CHQ = 0.004). There are several factors that likely contribute to this result. While the RfD for halofenozide (38 µg/kg/day) is higher than those for chlorpyrifos and cyfluthrin (5 and 25 µg/kg/day, respectively), its application rate was 1.7-fold higher than that for chlorpyrifos and 12-fold higher than that for cyfluthrin.

Since halofenozide has no USEPA assigned RfD, it has been calculated by Murphy and Haith (7). The authors used a conservative UF of 1000 due to the paucity of toxicological data. The halofenozide UF is 100-fold greater than chlorpyrifos (10) and 10-fold higher than cyfluthrin (100). This increased uncertainty also contributes to the relatively high CHQ. Also, there is no established dermal penetration factor for halofenozide, so a default value of 20% was used. This default value most likely overestimates the dermal absorption of halofenozide, and therefore the dose.

One factor that contributes to the relatively high CHQ is the dissipation of the halofenozide DFRs. Unlike azoxystrobin and carfentrazone-ethyl (and all other pesticides we have studied)




Figure 11. The distribution of halofenozide on whole body dosimeters. Hands, upper sock, and lower leg received the majority of the dose.

where there were significant reductions in available residues (43% and 37%, respectively) in the first hour following application, halofenozide DFRs were essentially unchanged over this time.

It has been postulated that pesticide residues on turf may become unavailable through absorption into the waxy layer of the leaf cuticle. Studies have shown that halofenozide penetrates insect cuticles poorly (11). Only 20% of the applied halofenozide penetrated the insect cuticle, compared to 45% for diflubenzuron and 85% for flucycloxuron, two other insect growth regulators. While insect and plant cuticles are different, both have a similar waxy layer at their outermost surfaces. It is possible, therefore, that the highly water soluble halofenozide also penetrates the plant cuticle very slowly.

The high water solubility may also influence the availability of halofenozide in other

ways. Halofenozide was watered in with 0.25 inches of irrigation like the other insecticides, so the thatch and verdure were saturated with water. The relatively high water solubility and low Koc (sorption to organic carbon) may mean that more of the halofenozide is in the water phase than bound to the thatch, increasing its availability during sampling. Halofenozide has also shown less vertical movement in bentgrass than in either bare soil or tall fescue (8). This reduced vertical movement may also increase the availability of halofenozide.

The reduction of available residues in the first hour for halofenozide (43%) and carfentrazone-ethyl (37%) is consistent with the ~ 50% reduction noted by Putnam and Clark (11) for a wide variety of pesticides. This new data further strengthens the case for a one-hour re-entry interval for treated turfgrass areas. The halofenozide

results over the first hour following application are unique. Halofenozide was the only compound tested in all the turf pesticides we evaluated that did not show a significant decline in available residues in the first hour following application. This finding shows that the reduction in residues in the first hour post-application may be widespread, but it is not universal.

The hands were determined to be the principle route of exposure for azoxystrobin (56%), carfentrazone-ethyl (28%), and halofenozide (54%). This route of exposure is in keeping with previous results for MCP, 2,4-D, and chlorothalonil, which, like azoxystrobin and carfentrazone-ethyl, received no post-application irrigation. The results for halofenozide are contrary to all other results found for insecticides that received post-application irrigation (4). The principle route of exposure for these compounds that received post-application irrigation was the lower body. Putnam and Clark (11) determined that 43-65% of the chlorpyrifos residues detected on the suit were found on the pants (15%), lower legs (20-35%), and socks (8-15%).

It is possible that the pesticides that did not receive post-application irrigation were sitting on the surface of the leaf and readily available to be picked up by casual hand contact. Following post-application irrigation, it is possible that while the residues could be picked up by the weighted California roller, they had moved past the verdure and were on/in the thatch and therefore unavailable for casual hand contact.

The high level of lower leg exposure for insecticides that received post-application irrigation may be the result of these compounds remaining in the water phase in the saturated thatch and verdure. Even one hour after the end of irrigation, the playing surface was wet. As the volunteer golfers walked across the turf, they may have been kicking up some water containing the pesticide applied thereby contaminating their lower body. It is possible that due to the relatively high water solubility of halofenozide (180-fold greater than chlorpyrifos, 36,000-fold greater than cyfluthrin) there was a much greater concentration of halofenozide in the water on the surface of the

thatch, leading to more availability to the hand.

The results from this study indicate that golfer exposure to turfgrass pesticides has a wide safety margin. It has also been shown that in two of the three cases, the use of reduced risk pesticides can substantially lower golfer hazard. Excluding halofenozide, it is also clear that a one-hour re-entry interval can also significantly reduce golfer hazard. In the cases examined, the use of reduced-risk pesticides can lessen golfer exposure and hazard. Azoxystrobin exposure following a simulated round of golf commencing 1 hour after application resulted in a CHQ (0.0039), >250-fold below current levels of concern.

Exposures to carfentrazone-ethyl (CHQ=0.00014, ~7,000-fold below) and halofenozide (CHQ=0.030, ~33-fold below) were also well under the USEPAs current level of concern. The application of azoxystrobin brought about a ~10-fold reduction in CHQ compared to chlorothalonil. The use of carfentrazone-ethyl produced an ~7,000 fold reduction in CHQ versus the use of MCP, and the application of halofenozide led to a ~2-fold reduction in CHQ compared to the use of chlorpyrifos.

While the use of halofenozide did not mitigate hazard as effectively as azoxystrobin and carfentrazone-ethyl, it is important to note that the hazard quotients calculated for all three reduced risk compounds are well below the USEPAs level of concern (1.0). Two of the three, azoxystrobin and carfentrazone-ethyl, can substantially reduce an already low risk. It is possible that the exposure profile for halofenozide was a factor in the loss of reduced-risk status for this compound.

Acknowledgements

The authors would like to thank the USGA's Turfgrass and Environmental Research Program for funding this work, and to Bayer Corporation for materials and technical assistance, and to Lisa Leombruni, Erin Shea, Barbara DeFlorio, and Alice Lao for their assistance in the laboratory and to all the volunteer golfers for their help in this project.

Literature Cited

1. Bernard, C. E., H. Nuygen, D. Truong, and R. I. Krieger. 2001. Environmental residues and bio-monitoring estimates of human insecticide exposure from treated residential turf. *Arch. Environ Contam. Toxicol.* 41(2):237-40. (TGIF Record 98498)
2. Chebira, S., Soltani, N., Muylle, S. and G. Smagghe. 2006. Uptake and distribution of three insect growth regulators – diflubenzuron, flucycloxuron and halofenozide – in pupae and adults of *Tenebrio molitor*. *Phytoparasitica* 34(2):187-196.
3. Cisar, J. L., R. H. Snyder, J. B. Sartain, and C. J. Borgert. 2002. Dislodgeable residues of chlorpyrifos and isazofos and implications for golfer exposure. *USGA Turfgrass and Environmental Research Online* 1(13):1-10. (TGIF Record 82908)
4. Clark, J. M., and J. Doherty. 2006. Managing pesticide exposure from treated turf. Page 35. In J. L. Nus (ed.). 2005 USGA Turfgrass and Environmental Research Summary. USGA, Far Hills, NJ. (TGIF Record 110124)
5. Fuller, R., D. Klonne, L. Rosenheck, D. Eberhart, J. Worgan, and J. Ross. 2001. Modified California roller for measuring transferable residues on treated turfgrass. *Bull. Environ Contam. Toxicol.* 67(6):787-794. (TGIF Record 78179)
6. Gardner, D.S. and B.E. Branham. 2001. Mobility and dissipation of ethofumesate and halofenozide in turfgrass and bare soil. *J. Agric. Food Chem.* 49:2894-2898. (TGIF Record 81576)
7. Murphy, R. R., and D. A. Haith. 2007. Inhalation health risk to golfers from turfgrass pesticides at three northeastern U.S. sites. *Environ. Sci. Technol.* 41(3):1038-1043. (TGIF Record 122901)
8. Occupational Safety and Health Administration. 1986. Method 62: chlorpyrifos, DDVP, diazinon, malathion and parathion in air. Organic Method Evaluation Branch, Occupational Safety and Health Administration, Analytical Laboratory Salt Lake City, Utah.
9. Pesticide manual, 14th edition. 2006. C.D.S. Tomlin, ed. British Crop Protection Council, Hampshire, UK.
10. Putnam, R. A., J. J. Doherty, and J. M. Clark. 2008. Golfer exposure to chlorpyrifos and carbaryl following application to turfgrass. *J. Agric. Food Chem.* 56:6616-6622. (TGIF Record 137962)
11. Putnam, R. A., and J. M. Clark. 2004. Managing pesticide exposure from turfgrass. *USGA Turfgrass and Environmental Research Online* 3(21):1-17. (TGIF Record 98807)