



TURFGRASS AND ENVIRONMENTAL RESEARCH ONLINE

USING SCIENCE TO BENEFIT GOLF



Runoff studies were carried out to identify which core cultivation practice, solid tine or hollow tine, maximized pesticide retention at the site of application. Measured quantities of pesticides in the edge-of-turf runoff and characteristics of a local golf course were used to calculate pesticide concentrations in surface water receiving turf runoff. Surface water concentrations of pesticides were compared to published toxicity data. Identifying management practices that reduce pesticide loss with runoff will improve disease and pest control in turf while minimizing undesirable environmental effects associated with the off-site transport of pesticides.

Volume 11, Number 8
August 1, 2012

PURPOSE

The purpose of *USGA Turfgrass and Environmental Research Online* is to effectively communicate the results of research projects funded under USGA's Turfgrass and Environmental Research Program to all who can benefit from such knowledge. Since 1921, the USGA has funded more than \$40 million for research at universities. The private, non-profit research program provides funding opportunities to university faculty interested in working on environmental and turf management problems affecting golf courses. The outstanding playing conditions of today's golf courses are a direct result of ***using science to benefit golf.***

Editor

Jeff Nus, Ph.D.
1032 Rogers Place
Lawrence, KS 66049
jnus@usga.org
(785) 832-2300
(785) 832-9265 (fax)

Research Director

Michael P. Kenna, Ph.D.
P.O. Box 2227
Stillwater, OK 74076
mkenna@usga.org
(405) 743-3900
(405) 743-3910 (fax)

Bill Katz, Chairman
Gene McClure, Co-chairman
Ed Michaels, Co-chairman
Ron Dodson
Kimberly Erusha, Ph.D.
Michael Fidanza, Ph.D.
Ali Harivandi, Ph.D.
Michael P. Kenna, Ph.D.
James Moore
Jeff Nus, Ph.D.
Paul Rieke, Ph.D.
James T. Snow
Clark Throssell, Ph.D.
Scott Warnke, Ph.D.
Chris Williamson, Ph.D.

Permission to reproduce articles or material in the *USGA Turfgrass and Environmental Research Online* (ISSN 1541-0277) is granted to newspapers, periodicals, and educational institutions (unless specifically noted otherwise). Credit must be given to the author(s), the article title, and *USGA Turfgrass and Environmental Research Online* including issue and number. Copyright protection must be afforded. To reprint material in other media, written permission must be obtained from the USGA. In any case, neither articles nor other material may be copied or used for any advertising, promotion, or commercial purposes.

Evaluation of Core Cultivation Practices to Reduce Ecological Risk of Pesticides in Runoff from Turf

Pamela Rice, Brian Horgan, and Jennifer Rittenhouse

SUMMARY

Pesticides associated with the turfgrass industry have been detected in stormwater runoff and surface waters of urban watersheds, raising concern of their potential environmental effects and a desire to reduce their transport to non-target locations. Runoff studies were conducted to compare the effectiveness of solid-tine versus hollow-tine core cultivation to reduce the quantity of pesticides transported with runoff from creeping bentgrass (*Agrostis stolonifera*) turf managed as a golf course fairway. The concentration of pesticides anticipated in a surface water receiving the runoff were calculated using data from this study and runoff volumes and pond dimensions recorded from a local golf course. Surface water concentrations were compared with levels known to be harmful to aquatic organisms. Key observations of the study were:

- Runoff volumes were less from turf managed with hollow-tine compared to solid-tine core cultivation.
- Greater quantities of pesticides were transported off-site with runoff from turf managed with solid tines.
- Concentrations of pesticides in a pond receiving runoff from turf managed with solid tines exceeded levels harmful to eight of 19 aquatic organisms evaluated.
- Replacing solid-tine with hollow-tine core cultivation reduced surface water pesticide concentrations to levels below harmful concentrations for most of these organisms.

Pesticides are applied to highly managed biotic systems such as golf courses, commercial landscapes, and agricultural crops. Golf course turf often requires multiple applications of pesticides at rates that exceed those typically found in agricultural or home environments (2, 12). Pesticides associated with the turfgrass industry have been detected in surface waters of urban

watersheds leading to increased concern about contaminant contributions from residential, urban, and recreational sources (6, 13, 15, 28, 32).

Fairways comprise approximately one-third of a typical golf course (29), which may border surface waters such as ponds, streams, and lakes. Golf course fairways and greens may be managed with core cultivation during the spring and/or fall to control thatch, alleviate surface compaction, enhance water infiltration, and stimulate root and shoot growth (3, 4, 5, 9, 27, 31). Cultivation with hollow tines typically involves removing cores from the turf, which are air-dried, broken down, and brushed back into the open holes. Solid-tine core cultivation requires a reduced amount of labor and is less disruptive to the surface of the turf, but it is believed to cause localized compaction (20).

Management practices have been shown to reduce runoff and pesticides transported with runoff from agricultural crops (14, 21, 24). Research on turfgrass also has shown the influence of cultural and irrigation practices on nutrient and pesticide transport with runoff and leachate (7, 10, 11, 16, 19, 25, 26, 30). The goal of this study was to identify which core cultivation practice, solid-tine or hollow-tine, maximizes pesticide retention at the site of application thus improving disease and pest control in turf while minimizing undesirable environmental effects associated with the off-site transport of pesticides.

Materials and Methods

Runoff Study Site

Experiments were conducted on turf plots managed as a golf course fairway at the University of Minnesota, Saint Paul, MN. The site (Waukegan silt loam) was divided into 6 plots

PAMELA RICE, Ph.D., Research Chemist, U.S. Department of Agriculture, Agricultural Research Service, Adjunct Professor, Department of Soil, Water, and Climate; BRIAN HORGAN, Ph.D., Associate Professor and Turfgrass Extension Specialist, Department of Horticulture; JENNIFER RITTENHOUSE, Physical Science Technician, U.S. Department of Agriculture, Agricultural Research Service; University of Minnesota, St. Paul, MN.



Figure 1. Creeping bentgrass turf managed with solid tine (A) or hollow tine (B) core cultivation. Cores removed with the hollow tines were air dried and worked back into the turf prior to pesticide application and simulated precipitation.

(24.4 m x 6.1 m, length x width) and sodded with ‘L-93’ creeping bentgrass (*Agrostis stolonifera*) sod 14 months prior to initiation of the reported studies. The turf was managed as a fairway with 1.25 cm height of cut (three times weekly, clippings removed), toppedressed with sand (weekly, 1.6 mm depth) and maintained with sprinkler irrigation.

Runoff collection systems were constructed at the western end of each plot, modified from the design of Cole et al. (7). Water traveled from the runoff gutter to a stainless steel flume equipped with an automated sampler and flow

meter. Gutter covers and flume shields prevented dilution of runoff with precipitation. Plots were hydrologically isolated with removable berms.

Management Practices

Plots were aerated twice (June 21, September 28) with either solid tines (0.95-cm diameter x 11.43-cm length with 5 cm x 5 cm spacing) or hollow tines (0.95-cm internal diameter x 11.43-cm length with 5 cm x 5 cm spacing) and toppedressed weekly with sand (Figure 1). Cores removed with the hollow tines were



Figure 2. A rainfall simulator delivered precipitation resembling storm intensities recorded in Minnesota. Runoff collection gutters guided runoff from the turf to flumes equipped with automated samplers and flow meters. Gutter covers and flume shields prevented dilution of runoff with precipitation.



Figure 3. A commercially available insecticide, fungicide, and herbicide were tank mixed and applied at label rates to all plots perpendicular to runoff flow; 63 days and 2 days following core cultivation and 26 (\pm 13) hours prior to initiation of simulated precipitation and runoff.

allowed to dry, broken into smaller pieces, and worked back into the turf. A backpack blower and leaf rake removed the turf and thatch from the plot surface. Sand topdressing was not performed immediately after core cultivation or within a week of simulated precipitation and generation of runoff.

Pesticide Application and Simulated Precipitation

A rainfall simulator was constructed to deliver precipitation similar to natural rain (8, Figure 2). Measured rainfall rates were similar to storm intensities recorded in Minnesota during July through October. Forty-eight hours prior to initiation of simulated precipitation each plot was pre-wet with the maintenance irrigation beyond soil saturation to allow for collection of background samples and to ensure uniform water distribution. Irrigation water samples and resulting background runoff were collected for analysis.

Petri dishes were distributed across the plots to verify pesticide application rates and rain gauges were distributed throughout each plot to quantify simulated precipitation. A commercially available insecticide, fungicide, and herbicide containing chlorpyrifos, flutolanil, mecoprop-p, dicamba, and 2,4-D were tank mixed and applied at label rates to all plots perpendicular to runoff flow (Figure 3). Simulated precipitation was initiated 26 (\pm 13) hours after pesticide application. Soil moistures were 46 (\pm 7)% water holding capacity within 3 hours prior to initiation of the simulated precipitation.

Runoff Collection and Pesticide Analysis

Runoff water samples were collected using automated samplers equipped with a flow meter to record water level in the flume, calculate flow rates, and total runoff volume. Time-paced water samples were removed from the automated samplers and stored frozen until laboratory analy-




Figure 4. Mean percentage of applied precipitation measured as runoff from turf plots managed with solid tine core cultivation or hollow tine core cultivation 63 days and 2 days prior to simulated precipitation and runoff. Error bars represent the standard deviation of the mean.

sis. Concentrations of chlorpyrifos, dicamba, flutolanil, MCPP, and 2,4-D were measured by direct injection of filtered samples onto a high performance liquid chromatograph with a photodiode array detector and quantified by direct comparison with external standard calibration curves of the analytical standards.

Calculating Pesticide Concentrations in a Pond Receiving Turf Runoff

Pesticide loads (micrograms m^{-2}) in the edge-of-plot runoff were calculated from recorded runoff volumes ($L m^{-2}$) and measured concentrations (micrograms L^{-1}) of pesticides in the runoff. Pesticide concentrations in a body of water receiving the runoff was determined using characteristics of a golf course located less than 20 miles from our study site, including the volume (L) of a pond receiving runoff from a known area of the golf course (m^2).

Estimated pesticide concentrations in a pond receiving runoff from fairway turf managed with solid-tine or hollow-tine core cultivation

were compared to published toxicity data to evaluate which core cultivation practice would be the most efficient at reducing environmental impacts. A detailed description of the calculations, toxicity data, and statistical analysis are provided elsewhere (23).

Results

Reduced Runoff Volume with Hollow-Tine Core Cultivation

Runoff volumes were reduced in fairway turf plots aerated with hollow-tine compared to solid-tine core cultivation. Although the period of time between core cultivation and simulated precipitation was greater for the first runoff event (63 days) than the second runoff event (2 days), due to a delay in the construction of the rainfall simulator, the overall trends observed between solid-tine and hollow-tine core cultivation remained the same, showing reduced runoff volumes with hollow tines for more than 80% of the samples (63 days = 81%, 2 days = 87%).




Figure 5. Chemographs and cumulative loads of chlorpyrifos (A) and 2,4-dichlorophenoxyacetic acid (2,4-D) (B) measured in runoff from turf plots managed with solid tines or hollow tines 2 days prior to simulated precipitation and runoff. Data for dicamba, flutolanil, and mecoprop-p (MCP) at 2 days and all pesticides at 63 days are provided elsewhere (23).

Calculation of cumulative runoff volumes from plots receiving core cultivation 63 days prior to rainfall simulation demonstrated a 10% reduction in cumulative runoff volume with hollow-tine compared to solid-tine cultivation. A 55% reduction in cumulative runoff volume with hollow-tine compared to solid-tine core cultivation was observed when plot received core cultivation 2 days prior to rainfall simulation.

The percentage of precipitation resulting as runoff from plots aerated with hollow tines was less than quantities observed from the solid-tine plots, suggesting greater infiltration with hollow-tine core cultivation (Figure 4). Other researchers have measured enhanced water infiltration in turf managed with hollow-tine core cultivation compare to untreated turf (1, 18) and greater saturated water conductivity and air porosity in turf managed with hollow tines compared to solid tines (20). The greatest difference in soil physical properties between plots was most prominent shortly after cultivation and diminishes with time as roots grow, compaction dissipates, and holes are covered or filled resulting in the greater distinction in runoff volumes between treatments at 2 days following cultivation compared to 63 days.

Reduced Pesticide Transport in Runoff with Hollow-Tine Core Cultivation

The quantity of pesticides transported with runoff from solid-tine plots exceeded that of the hollow-tine plots. Plots receiving hollow-tine core cultivation to manage thatch 63 days prior to runoff showed a 17, 15, 24 and 23% reduction in cumulative dicamba, flutolanil, MCPP and 2,4-D loads, respectively. Cumulative loads of chlorpyrifos were similar. Following the second core cultivation (2 days), hollow-tine plots displayed an even greater reduction in cumulative pesticide loads compared to the solid-tine plots with 46, 55, 37, 35 and 57% decline in cumulative loads of dicamba, flutolanil, MCPP, 2,4-D, and chlorpyrifos (Figure 5).

Correlation analysis of pesticide loads with runoff volumes and pesticide concentrations revealed pesticide loads were attributed to runoff

volume more than chemical concentrations for both management practices (volume $r = 0.78$ to 0.90 , concentration $r = 0.05$ to 0.22). This greater correlation of pesticide load with runoff volume explains in part the increased pesticide transport associated with the solid-tine plots compared to hollow-tine plots and the increased difference in pesticide loads between cultivation practices at 2 days compared to 63 days.

Hollow-tine core cultivation removed the cores and returned the soil back to the turf while solid-tine core cultivation pushed the soil aside to create the channels. As a result, one would anticipate greater soil compaction with the solid-tine cultivation and increased accessibility of soil adsorptive sites with the hollow-tine cultivation. This would influence hydraulic conductivity and infiltration as previously reported (1, 18, 20), as well as pesticide availability for transport (11, 17, 22).

The percentage of applied pesticides observed in the runoff is also influenced by the physical and chemical properties of the active ingredient. Chemical degradation was not influential in the present study as the time from chemical application to runoff (30 ± 8 hours) was much less than the reported half lives of the compounds of interest (5 to 320 days).

Reduced Risk of Pesticides in Receiving Surface Waters with Hollow-Tine Core Cultivation

Calculated concentrations of pesticides in a pond receiving runoff from fairway turf managed with hollow tines or solid tines were compared with published toxicological endpoints for 19 aquatic organisms including fish, amphibians, mollusks, crustaceans, aquatic plants, and algae (23). Toxicological endpoints included the median lethal concentration (LC_{50}) and median effective concentration (EC_{50} , i.e. the concentration of a compound that results in the measured effect in 50% of the organisms during a defined exposure period).

Pesticide levels in a surface water receiving runoff from turf managed with solid tines




Figure 6. Comparing concentrations of chlorpyrifos (A) and 2,4-dichlorophenoxyacetic acid (2,4-D) (B) in a surface water receiving runoff from fairway turf managed using solid tines or hollow tines 2 days prior to runoff with toxicological end points (median lethal concentrations or median effective concentrations) of sensitive aquatic organisms. Toxicity data available at http://cfpub.epa.gov/ecotox/ecotox_home.cfm.

exceeded the LC₅₀s or EC₅₀s for eight of the 19 evaluated aquatic organisms. With few exceptions, replacing solid-tine core cultivation with hollow-tine core cultivation reduced surface water concentrations of chlorpyrifos to levels below the LC₅₀ or EC₅₀ for three fish (Figure 6A), MCPP to levels below the EC₅₀ of a diatom (not shown), and 2,4-D to levels below the EC₅₀ of an aquatic plant (Figure 6B).

The sensitivity of rainbow trout, opossum shrimp, and water fleas to chlorpyrifos and water fleas to 2,4-D was great enough that surface water levels exceeded the LC₅₀s or EC₅₀s regardless of the turf cultivation practice (Figure 6A&B). Likewise, changes in management practice did not significantly influence the risk of pesticides to non-sensitive organisms (e.g. organisms whose LC₅₀ is well above the maximum concentration estimated in the diluted surface water. Data not shown).

Results of the present research provide quantitative information that will allow informed decisions on cultural practices that can maximize pesticide retention at the site of application, improving pest control in turf while minimizing environmental contamination and adverse effects associated with the off-site transport of pesticides. Using cultural practices that enhance infiltration and reduce runoff volume will effectively reduce pesticide runoff as demonstrated through the use of hollow-tine aeration.

Acknowledgements

The present research was funded, in part, by the USGA Turfgrass and Environmental Research Program. We thank Christina Borgen, Troy Carson, Mike Dolan, Shari Greseth, Andrew Hollman, Craig Krueger, Jeff Lanners, Jon Sass, Alex Seeley, Karli Swenson, and Nelson Irrigation.

Literature Cited

1. Baldwin, C. M., H. Liu, and P. J. Brown. 2006. Effects of core cultivation tine entry angle on golf putting greens. *HortTechnology* 16:265-269. (TGIF Record 203369)
2. Barbash, J. E., and E. A. Resek. 1996. Pesticides in groundwater: Distribution, trends, and governing factors. Lewis, Chelsea, MI. (TGIF Record 203412)
3. Beard, J. B. 1973. Turfgrass: science and culture. Prentice-Hall, Engelwood Cliffs, NJ. (TGIF Record 294)
4. Callahan, L. L., W. L. Sanders, J. M. Parham, C. A. Harper, L. D. Lester, and E. R. McDonald. 1998. Cultural and chemical controls of thatch and their influence on rootzone nutrients in a bentgrass green. *Crop Sci.* 38:181-187. (TGIF Record 42001)
5. Carrow, R. N., B. J. Johnson, and R. E. Burns. 1987. Thatch and quality of 'Tifway' bermudagrass turf in relation to fertility and cultivation. *Agron. J.* 79:524-530. (TGIF Record 10154)
6. Cohen, S., A. Svrjcek, T. Durborow, and N. L. Barnes. 1999. Water quality impacts by golf courses. *J. Environ. Qual.* 28:798-809. (TGIF Record 59340)
7. Cole, J. T., J. H. Baird, N. T. Basta, R. L. Huhnke, D. E. Storm, G. V. Johnson, M. E. Payton, M. D. Smolen, D. L. Martin, and J. C. Cole. 1997. Influence of buffers on pesticide and nutrient runoff from bermudagrass turf. *J. Environ. Qual.* 26:1589-1598. (TGIF Record 41754)
8. Coody, P. N., and L. J. Lawrence. 1994. Method and system for conducting meso-scale rainfall simulations and collecting runoff. U.S. Patent 5,279,151. Date issued: 18 January.
9. Dunn, J. H., D. D. Minner, B. F. Fresenburg, S.

- S. Bughrara, and C. H. Hohnstrater. 1995. Influence of core aeration, topdressing, and nitrogen on mat, roots, and quality of 'Meyer' zoysiagrass. *Agron. J.* 87:891-894. (TGIF Record 35964)
10. Evans, J. R., D. R. Edwards, S. R. Workman, and R. M. Williams. 1998. Response of runoff diazinon concentration to formulation and post-application irrigation. *Transactions of the ASAE* 41:1323-1329. (TGIF Record 203406)
11. Gardner, D. S., B. E. Branham, and D. W. Lickfeldt. 2000. Effect of turfgrass on soil mobility and dissipation of cyproconazole. *Crop Sci.* 40:1333-1339. (TGIF Record 69241)
12. Gianessi, L. P., and J. E. Anderson. 1995. Pesticide use in U. S. crop production. National Data Report. National Center for Food and Agricultural Policy, Washington, DC, USA. (TGIF Record 203404)
13. Gilliom, R. J., J. E. Barbash, C. G. Crawford, P. A. Hamilton, J. D. Martin, N. Nakagaki, L. J. Nowell, J. C. Scott, E. Stackelberg, G. P. Thelin, and D. M. Wolock. 2006. Pesticides in the nation's streams and ground water, 1992-2001. Circular no. 1291. Revised Feb 2007. U.S. Geological Survey National Water-Quality Assessment Program, Reston, VA, USA. (TGIF Record 203394)
14. Hansen, N. C., J. F. Moncrief, S. C. Gupta, P. D. Capel, and A. E. Olness. Herbicide banding and tillage system interactions on runoff losses of alachlor and cyanazine. *J. Environ. Qual.* 30:2120-2126.
15. Hoffman, R. S., P. D. Capel, and S. J. Larson. 2000. Comparison of pesticides in eight U.S. urban streams. *Environ. Toxicol. Chem.* 19:2249-2258. (TGIF Record 94952)
16. Kauffman, G. L., III, and T. L. Watschke. 2007. Phosphorus and sediment in runoff after core cultivation of creeping bentgrass and perennial ryegrass turfs. *Agron. J.* 99:141-147. (TGIF Record 120201)
17. Liu, L. X., T. Hsiang, and J. L. Eggens. 1995. Core cultivation and efficacy of benomyl applied to creeping bentgrass. *Agron. J.* 87:272-275. (TGIF Record 37385)
18. McCarty, L. B., M. F. Gregg, and J. E. Toler. 2007. Thatch and mat management in an established creeping bentgrass golf green. *Agron. J.* 99:1530-1537. (TGIF Record 131334)
19. Moss, J. Q., G. E. Bell, D. L. Martin, and M. E. Payton. 2007. Nutrient runoff from bermudagrass golf course fairways after aeration. *J. Appl. Turfgrass Sci.* (<http://dx.doi.org/10.1094/ATS-2007-0125-02-RS>). (TGIF Record 120171)
20. Murphy, J. A., P. E. Rieke, and A. E. Erickson. 1993. Core cultivation of a putting green with hollow and solid tines. *Agron. J.* 85:1-9. (TGIF Record 26844)
21. Potter, T. L., C. C. Truman, D. D. Bosch, and C. Bednarz. 2004. Flumeturon and pendimethalin runoff from strip and conventionally tilled cotton in the southern Atlantic Coastal Plain. *J. Environ. Qual.* 33:2122-2131.
22. Raturi, S., K. R. Islam, M. J. Carroll, and R. L. Hill. 2005. Carbaryl, 2,4-D, and triclopyr adsorption in thatch-soil ecosystems. *Environ. Sci. Health Part B* 40:697-710. (TGIF Record 203429)
23. Rice, P. J., B. P. Horgan, and J. L. Rittenhouse. 2010. Evaluation of core cultivation practices to reduce ecological risk of pesticides in runoff from *Agrostis palustris*. *Environ. Toxicol. Chem.* 29:1215-1223. (TGIF Record 163674)
24. Rice, P. J., J. A. Harman-Fetcho, A. M. Sadeghi, L. L. McConnell, C. B. Coffman, J. R. Teasdale, A. A. Abdul-Baki, J. L. Starr, G. W. McCarty, R. R. Herbert, and C. J. Hapeman. 2007. Reducing insecticide and fungicide loads in runoff

from plastic mulch with vegetative-covered furrows. *J. Agric. Food Chem.* 55:1377-1384. (TGIF Record 203368)

25. Smith, A. E., and D. C. Bridges. 1996. Movement of certain herbicides following application to simulated golf course greens and fairways. *Crop Sci.* 36:1439-1445. (TGIF Record 39465)

26. Starrett, S. K., N. E. Christians, and T. A. Austin. 1996. Movement of pesticides under two irrigation regimes applied to turfgrass. *J. Environ. Qual.* 25:566-571. (TGIF Record 37607)

27.. Turgeon, A. J. 1985. Turfgrass management, 2nd ed. Reston Publishing, Reston, VA. (TGIF Record 13745)

28. USEPA. 1999. National recommended water quality criteria. USEPA 822-Z-99-001. United States Environmental Protection Agency, Washington, DC, USA.

29. Watson, J. R., H. E. Kaerwer, and D. P. Martin. 1992. The turfgrass industry. Pages 29-88. In D. V. Waddington (ed.). Turfgrass. Agron. Monogr. 32. ASA, Madison, WI. (TGIF Record 26020)

30. Wauchope, R. D., R. G. Williams, and L. R. Marti. 1990. Runoff of sulfometuron-methyl and cyanazine from small plots: Effects of formulation and grass cover. *J. Environ. Qual.* 19:119-125. (TGIF Record 16977)

31. White, R. H, and R. Dickens. 1984. Thatch accumulation in bermudagrass as influenced by cultural practices. *Agron. J.* 76:19-22. (TGIF Record 458)

32. Wotzka, P. J., J. Lee, P. D. Capel, and M. Lin. 1994. Pesticide concentrations and fluxes in an urban watershed. Pages 135-145. In AWRA Technical Publication Series TPS-94-4. American Water Resources Association, Herndon, VA, USA, (TGIF Record 203372)